

On the Possible Sources of D/H Dispersion at High Redshift

Brian D. Fields

Department of Astronomy, University of Illinois, Urbana, IL 61801, USA

Keith A. Olive

*Theoretical Physics Institute, School of Physics and Astronomy,
University of Minnesota, Minneapolis, MN 55455 USA*

Joseph Silk

*Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH,
and Institut d'Astrophysique, 98 bis Boulevard Arago, Paris 75014, France*

Michel Cassé

*Service d'Astrophysique, CEA, Orme des Merisiers, 91191 Gif sur Yvette, France
also Institut d'Astrophysique, 98 bis Boulevard Arago, Paris 75014, France*

and

Elisabeth Vangioni-Flam

Institut d'Astrophysique, 98 bis Boulevard Arago, Paris 75014, France

ABSTRACT

Recent observations suggest the existence of a white dwarf population in the Galactic halo, while others suggest that deuterium has been astrated in systems at high redshift and low metallicity. We propose that these observations could be signatures of an early population of intermediate-mass stars. Such a population requires a Population III initial mass function different from that of the solar neighborhood, as perhaps also suggested by the observed cosmic infrared background. Also, to avoid overproduction of C and N, it is required that the $Z = 0$ yields of these stars have low ($\sim 10^{-3}$ solar) abundances as suggested by some recent calculations. Under these assumptions, we present a model which reproduces the observed D vs Si trend, and predicts a high cosmic Type Ia supernova rate, while producing a white dwarf population that accounts for only $\sim 1.5\%$ of the dark halo. This scenario can be tested by observations of the cosmic supernova rate, and by confirmation and further studies of the putative white dwarf halo population.

1. Introduction

One of the key parameters in the standard big bang model is the density of ordinary baryonic matter, expressed as a fraction of the critical density, Ω_B (often quoted in combination with the Hubble parameter as $\Omega_B h^2$). Big bang nucleosynthesis (BBN) has traditionally been the primary theoretical tool for determining $\Omega_B h^2$, and has recently been complemented by detailed measurements of the cosmic microwave background anisotropy. These results are to be compared to direct surveys of cosmic baryonic matter. While baryons are readily observable in stars or stellar remnants, galaxies, X-ray-emitting hot gas, as well as intergalactic gas, the low-redshift baryonic budget (Fukugita, Hogan, & Peebles (1998)) is uncertain, and leaves some room for dark baryonic objects. Conversely, at higher redshifts, ($z = 2$ to 3) a new estimate of $\Omega_B h^2$ based on the measurement of Lyman alpha forest intergalactic H I absorption lines (and also He II lines) in the spectra of remote quasars has been done and is in good agreement with $\Omega_B h^2$ derived from BBN (in the range $\Omega_B h^2 = 0.01$ to 0.03 , Riedeger *et al.* (1998), Wadsley *et al.* (2000)). This is between 3 and 20 times the observed baryon density in the luminous parts of galaxies and their presumed precursors, the damped Lyman alpha intergalactic neutral hydrogen clouds, amounting to $\Omega_B h^2 = 0.0015$ to 0.003 (Pérourx *et al.* 2001).

At the local scale, MACHOS, with masses of about $0.5 M_\odot$ and detected by gravitational microlensing towards the LMC, are possible candidates for some of the dark baryons, and amount to $\lesssim 20\%$ of the dark halo (Alcock *et al.* 2000). Recently, old halo white dwarfs may have been detected in sufficient numbers to amount to $\gtrsim 2\%$ of the dark halo. These white dwarfs, if halo members, represent a large increment, by a factor ~ 10 , relative to disk white dwarfs, and hence would have a profound impact on Galactic chemical evolution. In this paper we discuss the implications of the discovery of dark baryons in the form of old halo white dwarfs for BBN, and in particular for the deuterium abundance.

The standard model of BBN has enjoyed great success in its ability to predict the primordial abundances of the light element isotopes, D, ^3He , ^4He and ^7Li . Indeed, in the standard model, there remains only one remaining unknown parameter, the baryon-to-photon ratio, $\eta = n_B/n_\gamma$, which can be expressed as $\eta_{10} = 10^{10}\eta = 274\Omega_B h^2$. Roughly, for a value of η_{10} between 1.7 and 7.4, (corresponding to $\Omega_B h^2$ between 0.006 and 0.027), we obtain concordance between the BBN predictions and the observational determination of the abundances (Nollett & Burles 2000; Vangioni-Flam, Coc, & Cassé 2001; Cyburt, Fields, & Olive 2001). In detail, however, there are small but perhaps not insignificant differences in the determination of $\Omega_B h^2$ from the different isotopes. ^4He and ^7Li concordance is typically achieved for lower η_{10} whereas D/H is best fit at higher values. To be sure, not every isotope serves as an equally good baryometer. In fact, because of the weak logarithmic dependence on η_{10} ,

^4He is the worst element to use to fix the baryon density. While the ^7Li abundance strongly depends on η_{10} , the theoretical uncertainties which stem from uncertainties in the input nuclear cross-section make it difficult to derive precise values of the baryon density. In addition, because of the functional dependence of ^7Li on η_{10} , a given observational abundance of ^7Li gives us two predictions for η_{10} . D/H, on the other hand, is both strongly dependent on η_{10} and is relatively free from theoretical uncertainties. One should bear in mind however, that there are far more observational determinations of ^4He and ^7Li at the present time than there are of D/H from quasar absorption systems. Nevertheless it is the range in D/H, reported to span up to a factor of 20, that has been the center of much controversy over the past decade because of the strong sensitivity to the baryon density. The reported D/H abundance has now converged towards the lower end of this range, but recent measurements reveal a spread by up to a factor of 3 that may reflect intrinsic variations in the D/H abundance. These variations would most plausibly arise from stellar processing of primordial gas, or possibly from non thermal processes such as photodisintegration by gamma rays, or spallation reactions induced by energetic particles.

The recent results BOOMERanG yield $\Omega_B h^2 = 0.021^{+0.004}_{-0.003}$ (Netterfield *et al.* 2001), in very good agreement with the estimate from D/H. This is also in agreement with the recent results of DASI (Pryke *et al.* 2001), who find $\Omega_B h^2 = 0.022^{+0.004}_{-0.003}$.

There have also been several new observations of D/H from quasar absorption systems. In addition to the earlier determinations by Burles and Tytler (1998a,b) in two Lyman limit systems of $4.0 \pm .65$ and $(3.25 \pm 0.3) \times 10^{-5}$ and the upper limit in a third Lyman limit system by Kirkman *et al.* (1999) of 6.8×10^{-5} , there have been three new measurements of D/H in Damped Lyman α systems by O’Meara *et al.* (2001): 2.5 ± 0.25 , by D’Odorico *et al.* (2001): 2.25 ± 0.65 , and by Pettini and Bowen (2001): 1.65 ± 0.35 all times 10^{-5} . This last value is uncertain due to the poorly defined level of the continuum and the possible blending of lines (P. Petitjean, private communication). We note that there is also a measurement of D/H in a relatively low redshift system with significantly higher D/H (Webb *et al.* 1997; Tytler *et al.* 1999). This system has recently been criticized (Kirkman *et al.* 2001) and we will not consider it further here.

Is there a real dispersion in D/H in these high redshift systems? We note two possible trends that go in a similar direction. The data may show an inverse correlation of D/H abundance with Si. This may be an artifact of poorly determined Si abundances, or (as yet unknown) systematics affecting the D/H determination in high-column density (damped Lyman- α , hereafter DLA) or low-column density (Lyman limit systems) absorbers. On the other hand, if the dispersion in D/H is real, it has profound consequences, as it must indicate that *some processing of D/H must have occurred even at high redshift*. The second trend is

that the data may show an inverse correlation of D/H abundance with HI column density. If real, this would suggest that in the high column density DLA systems, which are most likely to have undergone some star formation, some processing of D/H must similarly have occurred, likewise at high redshift.

It is interesting to note that local D values measured in the interstellar medium also appear to show a considerable dispersion. Linsky (1998) summarized the observations then available and argued for a constant value in the local ISM of $(1.5 \pm 0.1) \times 10^{-5}$. Recent data (see Vidal-Madjar 2001, for a review) are consistent with this mean value, but suggest significant variations of the D/H ratio spanning the range 5×10^{-6} to 4×10^{-5} . Thus, if D shows such variability, perhaps correlated to some unexpected physical processes leading to its destruction, one can, in turn, question the ability of cosmological clouds to represent the genuine primordial abundance.

Thus, a simple average of D/H abundance determinations does not make sense, at least without a proper enlargement of the error in the mean due to the poor χ^2 that such a mean would produce. Moreover, if deuterium destruction has occurred, we must question the extent to which any of these systems determine the value of Ω_B .

There is one obvious objection to the hypothesis that nucleosynthetic processing might have occurred in the high-redshift systems. The absorbing systems have low metallicity. We now point to seemingly unrelated observations that indicate that primordial processing may have occurred, and with substantially less enrichment than is associated with standard yields for a solar neighborhood initial mass function (IMF).

A substantial population of low luminosity white dwarfs has recently been discovered (Oppenheimer *et al.* 2001; Ibata *et al.* 2000). These objects were identified by their large proper motion, indicating halo kinematics. Oppenheimer *et al.* find that if the objects are a halo population of white dwarfs, then they represent a fraction $\gtrsim 2.5\%$ of the dark halo mass. There has been some question about whether these objects do reside in the halo (Hansen 2001; Reid, Sahu and Hawley 2001), but a recent re-analysis of the data by Koopmans & Blandford (2001) also concludes that these are halo objects, and finds a similar mass budget to that of Oppenheimer *et al.* We now address the potential implications of any observed dispersion in D/H in high redshift quasar absorption systems for a relationship between the possibility of D/H destruction in these systems by intermediate mass stars and the ensuing production of a population of halo white dwarfs.

We summarize the importance of D as a baryometer, and its relation to the other light element abundances, in §2. In §3 we argue that the observed dispersion in high-redshift D/H, if real, cannot be accounted for by standard chemical evolution models. Thus, an explanation

of such a dispersion will necessarily involve nonstandard scenarios, involving, e.g., variations in the IMF, or outflows of material. We search for models which can explain this dispersion as well as the white dwarf observations. We find that such models can be constructed, assuming an initial burst of intermediate-mass star formation. It is also necessary that the C and N yields of zero metallicity stars are small, as suggested by some recent models. The prospects for testing this model and the implications for BBN are summarized in §4.

2. Determinations of Ω_B

As discussed above, each of the light elements can be individually used to determine $\Omega_B h^2$ with varying accuracy. For example, a helium mass fraction of $Y = 0.238$ corresponds to $\eta_{10} = 2.4$ or $\Omega_B h^2 = 0.009$. However, the systematic uncertainties in the ${}^4\text{He}$ abundance make it difficult to exclude values of Y as high as 0.25 (see for example, Olive & Skillman 2001) which because of the weak dependence on η corresponds to $\eta_{10} = 7.4$ or $\Omega_B h^2 = 0.027$. With ${}^7\text{Li}$ one can do much better. A plateau value of ${}^7\text{Li}/\text{H} = 1.2 \times 10^{-10}$ corresponds to $\eta_{10} = 2.4$ and 3.1. However, the slightly higher value of ${}^7\text{Li}/\text{H} = 2 \times 10^{-10}$ (consistent within systematic uncertainties) yields $\eta_{10} = 1.7$ and 4.5. While there are occasional attempts to further increase the ${}^7\text{Li}$ abundance on the basis that some ${}^7\text{Li}$ has been depleted (Vauclair & Charbonnel 1998; Pinsonneault *et al.* 1998,2001; Theado & Vauclair 2001), it is very hard to do so, due to recent ${}^6\text{Li}$ observations which leave little room, if any, for ${}^7\text{Li}$ depletion (Fields & Olive 1999; Vangioni-Flam *et al.* 1999) and the lack of dispersion in the Li data (Bonifacio & Molaro 1997; Ryan, Norris, & Beers 1999; Ryan *et al.* 2000).

In contrast, the strong dependence of D/H on η makes deuterium a very good baryometer, provided we have an accurate determination of D/H. The values of D/H measured in the high redshift systems however cover the range $\eta_{10} = 4.8$ for $\text{D}/\text{H} = 4 \times 10^{-5}$ to $\eta_{10} = 8.5$ for $\text{D}/\text{H} = 1.65 \times 10^{-5}$. Thus if the dispersion in the D/H is real (beyond observational error as the quoted error bars imply), then one can question the extent that any of these determinations truly represent the primordial D/H abundance.

In the event that the dispersion is real, some amount of D/H must have been destroyed, presumably by stellar processing. Since $\text{D}/\text{H}(t)$ is monotonically decreasing in time, the primordial value should be greater than or equal to the maximum of the observed values.

Stellar processing must involve short-lived stars of intermediate or even larger mass. The intermediate mass stars are the proposed white dwarf precursors. However, even an IMF peaked near $2\text{--}8 M_\odot$ could include some massive stars above $10 M_\odot$ that would explode as SNII and produce the heavy element enriched observed in the absorption systems. An

inverse correlation of D with Si abundance is a natural consequence. Since these enriched systems are preferentially the more massive and advanced with respect to disk formation, as we discuss below, they might also have higher gas column densities and account for the additional inverse correlation of D with HI column.

3. Destruction of D/H by galactic chemical evolution

The high column density quasar absorption systems are generally considered to be protogalactic disks. In particular DLAs have the kinematics and ionization characteristic of the interstellar medium of massive protodisks. Lyman limit systems have a greater extent and display higher ionization than the DLAs. Star formation with a normal IMF could not possibly have occurred in these systems to any substantial extent. Typical metallicities are 0.1 to 0.01 solar, and this is consistent with these systems being predominantly gaseous. However in order for there to be enough halo white dwarfs to account for the recent observations, the primordial IMF must have been very different from the present day IMF, and been dominated by intermediate mass stars. These stars would have formed during the protogalactic phase, and could have destroyed significant amounts of primordial deuterium.

Typical stellar D/H depletion mechanisms usually predict the production of heavy elements in closed box models. Indeed, by the time the metallicity reaches 0.1 solar abundance in a standard galactic evolution model, only a few percent of the initial D/H ratio has been affected. This is the main reason one is led to associate the observed D/H ratio to that of the primordial deuterium abundance. It is conceivable, however, that an early generation of stars was formed with masses only in the intermediate mass range of $2-8 M_{\odot}$. These stars would not be significant producers of heavy elements (oxygen and above) usually ascribed to Type II supernovae. Motivation for this suggestion comes from the discovery by Oppenheimer *et al.* (2001) of a population of old white dwarfs in the halo of the Galaxy (see also Ibata *et al.* 2000; Chabrier 1999; Gates & Gyuk 2001).

Previous studies of white dwarf halos have examined the consequences when the entire halo consists of white dwarfs. There are many objections to this extreme hypothesis, ranging from overproduction of He, excessive halo Type Ia supernova rates, excessive light element production in the early (proto-)Galaxy, and excessive C and N production. Constraints on the high-redshift population, and possible observable signatures have been studied by several groups (Ryu, Olive, & Silk 1990; Fields, Freese & Graff 2000; and references therein). The principal refinement we make to these ideas is to note that the old white dwarf halo

population is sub-dominant, possibly amounting to as little as 2 percent of the dark halo.¹ Nevertheless, the effects on galactic chemical evolution are profound.

To see this, let consider the mass budget for deuterium destruction. In order to achieve noticeable D depletion, one must process a significant amount of material through an early generation of stars. A typical galaxy may have $M_B = 10^{11} M_\odot$ of visible baryons, mostly in the form of stars, and about $10^{12} M_\odot$ total, the balance being dark matter. We point out that a white dwarf halo mass fraction of even 2% can have significant consequences for D destruction in protodisk and disk gas. This yields about $2 \times 10^{10} M_\odot$ in white dwarfs, and hence $\sim 10^{11} M_\odot$, comparable to the present disk mass, in precursor stars. All $8 \times 10^{10} M_\odot$ of the ejecta is totally astrated. Hence present-day astration factors of 2 or more are easily achievable with a white dwarf halo mass fraction of 1% or less, in an interstellar medium containing some $6 \times 10^9 M_\odot$ in atomic and molecular gas.

The requirements of D astration and white dwarf production, all with little heavy element production, lead to very strong constraints on the required IMF. To avoid heavy element production through Type II supernovae, we must demand that the IMF is restricted to the $2\text{--}8 M_\odot$ range (Adams & Laughlin (1996)). At the $m < 2 M_\odot$ end, the constraint is that too many stars would be luminous today or would have been seen as a huge population of red giants at $z \sim 3$, as $t \simeq 2$ Gyr at this redshift in an $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$ universe. On the $m > 8 M_\odot$ end, the problem is that we do not want to make a lot of metals (oxygen and up) from supernovae. This constraint is not as strong, and we need to add a few supernovae to get some metal production for the absorption systems.

A new result which independently demands an altered IMF at high redshift comes from the far IR background. Models of the FIRB (by ULIGs at high z) overproduce K light today (ie too many low mass stars) if a standard IMF is adopted (Cole *et al.* 2001). An IMF peaked at intermediate masses offers an attractive solution: it avoids the low mass stars and overproduction of heavy elements but gives the light. The FIRB requires νi_ν to be about equal to what is produced with a standard IMF for the optical/near-IR background, i.e., about $10 - 15 \text{ nW/m}^2$ (and to be produced in dust-shrouded ULIGS; see, e.g., Hauser and Dwek 2001 for a recent review). The white dwarfs that the ULIGS (which are known to be merger-induced bursts of star formation) produce would then (possibly after another merger or if they formed in outflow-boosted star formation) be the halo white dwarfs. The kinematics are a weak point but the moral is striking: a peaked IMF may be needed for

¹It has been argued (Reid *et al.* 2001)) that the detected white dwarfs might be a thick disk population. If this is the case, the total mass of white dwarfs is difficult to determine precisely but is considerably smaller and would thus not have important consequences for deuterium.

completely different reasons.

The obvious objection is that the hypothesised precursor stars will overproduce C and possibly N that would be observable in Galactic or extragalactic material (e.g., Fields, Freese, & Graff 2000). This is a serious issue due to the nature of the observed C and N abundances in primitive systems. For our purposes, the most constraining C and N values are those in the same QSO absorbers for which D has been detected. Namely, C has been detected in the best-measured Lyman limit systems (Burles & Tytler (1998a; 1998b); O’Meara *et al.* (2001)), and in all three cases $[C/H] \sim -2$, and $[C/O] \lesssim 0$ or $[C/Si] \lesssim 0$. Results for N are similar. Thus, the observations apparently demand that no significant C or N enrichment (over that of α -elements) takes place.

These observations place key constraints on a putative intermediate-mass Population III. If the yields of zero-metallicity intermediate mass stars are comparable to those of their low-metallicity counterparts (e.g., van den Hoek & Groenewegen (1997)), then indeed, the resulting enrichment is extremely large if most of the baryons are processed through white dwarfs. Although the enrichment is reduced when only a fraction of all baryons are processed, even here, to make a white dwarf halo at 10% of the total leads to C and N abundances at about $[C, N/H] = -1$. As this violates the observational constraints, we conclude that a halo white dwarf population cannot be created at high redshift if $Z = 0$ intermediate-mass stars create significant C and N. It then also follows that the dispersion in D due to chemical evolution is small; a similar conclusion was reached by Prantzos & Ishimaru (2001).

It is far from clear, however, that zero-metallicity intermediate-mass stars behave in this way. In fact, the nucleosynthetic yields for these stars are quite uncertain. In this mass range, at all metallicities, the yields depend sensitively on the details of several “dredge-up” events in which convective mixing brings nuclear burning products up to the envelope of the star. These have been studied in detail for solar metallicity stars and for moderately metal-poor stars, and the main results of different groups are in reasonable qualitative and quantitative agreement. However, there remain basic qualitative uncertainties in the post-main sequence evolution of zero-metallicity intermediate-mass stars.

Specifically, it is as yet unclear—i.e., model-dependent—whether the last and most important of these mixing events (the “third dredge-up”) occurs. There has been considerable effort recently to answer this question, but the studies have given conflicting results. Chieffi, Domínguez, Limongi, & Straniero (2001) do find such a phase, and conclude that zero-metallicity intermediate-mass stars should be important sources of C and N. The work of Marigo, Girardi, Chiosi, & Wood (2001) does not find a third dredge-up, but the authors are uncertain whether such a phase would have occurred had they run their models longer; what *is* clear is that the C and N envelope abundances *prior* to this phase are small ($\lesssim 10^{-3}$

of solar) and so would not create a significantly signature. Finally, Fujimoto, Ikeda, & Iben (2000) argue that for stars with $M \gtrsim 4M_\odot$ and $[\text{Fe}/\text{H}] \lesssim -5$, there is no third-dredge up at all.

It thus appears that the question of C and N abundances in $Z = 0$ stars is both complex and unsettled. To simultaneously explain a halo white dwarf population and D depletion at high redshift requires that these stars do not eject significant C and N. For the moment, this amounts to an assumption, but one with some support from ongoing studies.

To make the proposed scenario concrete and quantitative, we have constructed chemical evolution models which lead to significant D astration at low metallicity and also have significant white dwarf production. We will present one such model; of course, given the freedom to adjust the IMF (and star formation history), a large class of models are allowed. However even with this freedom, the acceptable models share common traits and make similar predictions. We have explored the case of a closed-box model. The stellar creation or birth function $C(m, t)$ encodes the rate and mass spectrum of new star birth via

$$dN_\star = C(m, t) dm dt \quad (1)$$

We adopt the “bimodal” form $C(m, t) = \psi_1(t)\phi_1(m) + \psi_2(t)\phi_2(m)$. We choose the star formation rate $\psi_1 = (M_1/\tau_1)e^{-t/\tau_1}$ to be a short-lived burst of star formation with $\tau_1 = 100$ Myr and $M_1 = 0.75M_B$, with an associated IMF which is sharply peaked at intermediate mass stars; this is the population which destroys deuterium and creates white dwarfs. Following Adams & Laughlin (1996), we take the IMF to be lognormal

$$\phi_1(m) \propto m^{-1} \exp[-\ln^2(m/m_c)/2\sigma^2] \quad (2)$$

The model we present here has a characteristic mass scale $m_c = 5M_\odot$ and a dimensionless width $\sigma = 0.07$ (this is very similar to a gaussian with mean $\mu = 5M_\odot$ and width $\sigma_G = \sigma m_c = 0.35M_\odot$). Other parameter choices are possible; for higher m_c , the needed σ is narrower. The other term in the creation function describes standard, quiescent star formation. We have $\psi_2 = \lambda g(t)M_{\text{gas}}$, where $\lambda = 0.3 \text{ Gyr}^{-1}$, and the factor $g(t) = 1 - e^{-t/0.5\text{Gyr}}$ allows for a smooth transition from the bursting to the quiescent star formation rate. The IMF ϕ_2 is a power law with slope 1.35 (Salpeter).

Results from the model appear in Figure 1. In panel (a) we see the evolution of D/H versus metallicity $[\text{O}/\text{H}]$ (as O is an α -element which is co-produced with Si, we take $[\text{O}/\text{H}] = [\text{Si}/\text{H}]$). Three models are plotted. The solid curve corresponds to a bimodal model in which we have taken a primordial value $\text{D}/\text{H} = 5.0 \times 10^{-5}$. The two phases of star formation are clear, as each leads to a characteristic drop in D, the burst phase acting at low metallicity and the quiescent phase acting at high metallicity. The points are the observed data discussed




Fig. 1.— Results from the bimodal star formation model. **(a)** D evolution as a function of metallicity. Data and model curves are as noted in the text; the protosolar value is that of Geiss & Gloeckler (1998); the ISM value and errorbar of Linsky (1998) appears as the solid line, while the range reported by Vidal-Madjar appears as the dotted errorbar. **(b)** D evolution as a function of remnant production as measured by the total Galactic mass M_{rem} in remnants, almost all of which are white dwarfs. **(c)** The predicted cosmic supernova rates, in arbitrary but comoving units, as a function of redshift.

earlier. The values from quasar absorption line systems have an arrow indicating the Si abundances are lower limits.² The dashed curve in Figure 1 corresponds to a bimodal model with $\text{D/H} = 4.0 \times 10^{-5}$. In this case, there is less astration needed at low metallicity, so the amount of gas processed in the Pop III burst is lower, i.e., $M_1 = 0.5M_B$ (as is the resulting mass of halo white dwarfs). Note that the two model curves join once the D has been destroyed in the Pop III phase, and follow the same quiescent evolution afterwards. We see that in both cases, the curves provide a reasonable fit to the data. Therefore, it is difficult to use the existing D/H data and chemical evolution to accurately predict the primordial

²This arises because the Si abundances are measured in the gas phase, and do not include any Si which has been depleted onto dust grains. The presence of grain depletion complicates attempts to interpret measurements and trends in high-redshift abundances, and to make matters worse, the depletion levels found in the Milky Way (more than an order of magnitude) are apparently not always appropriate for high-redshift systems. As shown for the case of DLAs (e.g., Pettini, Ellison, Stidel, Shapley, & Bowen 2000; Molaro *et al.* 2000; Prochaska & Wolfe 1999; Lauroesch, Truran, Welty, & York) Si/Zn can be much higher than in the ISM. Si/Zn is typically about 1/3 (i.e., 0.5 dex), and varies from system to system, sometimes exceeding the solar ratio.

D/H value and hence $\Omega_B h^2$. Finally, the dot-dashed curve corresponds to a model in which there is no Pop III and a primordial value $D/H = 3.0 \times 10^{-5}$ as in O’Meara et al. (2001). In this “standard” case there is essentially no D depletion at low metallicity and, as we have noted, the dispersion in D is unexplained.

The deaths of the Pop III intermediate-mass stars not only release D-free material, but also produce white dwarfs. This correlation is illustrated in Figure 1(b); we plot the D abundance versus the total mass in compact remnants, essentially all of which is in white dwarfs. The model used here is that shown in the solid curve of Figure 1(a). We see that the burst phase, during which D/H drops to about 2.1×10^{-5} , puts about 15% of the baryons (about $1.5 \times 10^{10} M_\odot$) into white dwarfs. The quiescent phase adds another $10^{10} M_\odot$. Note well the distinction between this work and previous work. Previous work was centered on the possibility of a white dwarf halo. Here we are interested in D/H and producing a few white dwarfs. A halo population of a few percent would dramatically ease the previous constraints. Indeed, Figure 1(b) shows that the 2.5% observation of Oppenheimer is more than sufficient.

Note that by showing the extragalactic D *vs.* Si data on the same plot as the Galactic points, we implicitly assume that the Galaxy is a system representative of the quasar absorbers. It then would also follow that the white dwarf and carbon production also would apply to these systems. However, one need not take the model so literally as to demand that all baryonic matter—or even all the baryons in protogalaxies—follow this evolution. For example, in a hierarchical merging picture, one could instead imagine that this evolution is common to some but not all of the subsystems which form the Galaxy. Thus, D depletion (and carbon and white dwarf production) would be inhomogeneous within the protogalaxy, and across the observed quasar absorbers.

Figure 1(c) shows the predicted supernova rates, both for Type II as well as Type Ia events. The model use here is that shown in the solid curve of Figure 1(a) and in Figure 2. The Type II rate follows the assumed star formation rate, which has simply been taken to be $\psi \propto M_{\text{gas}}$ and which turns on sequentially after the bursting Pop III star formation ends. The Type Ia rate is calculated using the prescription of Matteucci & Greggio (1986), and assumes that the binarity properties of the Pop III stars are the same as those observed locally today. The Type Ia rate peaks sharply, echoing the rapid burst and small mass range that characterized the Pop III star formation. Note that while the peaked nature of the Ia rate is a generic feature of the model we propose, the peak redshift is model-dependent, reflecting the characterist mass (and thus the lifetime) of the Pop III stars. Figure 1(c) shows the results for a value $m_c = 5M_\odot$, which gives a Type Ia peak at $z \sim 3$. Had we chosen $m_c = 6M_\odot$ ($4M_\odot$), we would find a peak at $z \sim 5$ (2). In any case, we predict that the Type Ia supernova rate is quite large at moderate redshift; this can be tested in

ongoing high- z supernova searches. Note that a suppression of Type Ia supernovae at low metallicities (Kobayashi *et al.* (1998)), if real, could alter this prediction.

A potential constraint on high-redshift Type Ia supernova activity also arises from the cosmic γ -ray background at MeV range. The dominant source of photons at these energies is mostly nuclear line emission due to the $^{56}\text{Ni} \rightarrow ^{56}\text{Co} \rightarrow ^{56}\text{Fe}$ decay chain (e.g., Ruiz-Lapuente, Cassé, & Vangioni-Flam (2001)). The enhanced SN Ia production of our model increases the MeV background relative to the predictions expected from a constant-IMF cosmic star formation rate. However, most of the contribution is at high redshift and is thereby diluted. In fact, the background is not a good discriminant of these epochs, and the Type Ia rate proposed here is non inconsistent with the cosmic MeV background.

4. Conclusions

The present work is motivated by the reports of a population of halo white dwarfs in our Galaxy, and of dispersion in the high redshift D/H values. In light of these recent observations, we have explored the consequences of a high redshift population of intermediate mass stars. This population leads to D destruction at low metallicity, as well as white dwarf production, and would add to the far-IR background without increasing the local near-IR background. For this scenario to remain viable, it is necessary that the C and N yields of zero-metallicity stars are small, as suggested by some recent stellar evolution models. We have constructed relatively simple models which fit the data reasonably well for primordial abundances of $\text{D}/\text{H} = 4$ and 5×10^{-5} . Other values are possible by merely adjusting the astration constant, M_1 , in the POP III star formation rate. Thus, we conclude that it is difficult to use existing data to accurately predict either the primordial abundance of D/H or $\Omega_B h^2$.

An interesting prediction of a halo in which the white dwarf number exceeds that in the disk by an order of magnitude is that, binarity fractions being comparable, the number of Type Ia would be greatly boosted in the very early universe. Some may even be seen in the halos of nearby galaxies, (Smecker & Wyse 1991), though the spatial distribution of these events within galaxies will depend on the interplay between the formation and evolution of the Pop III stars and protogalaxies.

A potentially disturbing complication is raised by the work of Hansen (2001), who estimates the white dwarf ages from their luminosity and color distribution. He finds that they are relatively young and if true, this is inconsistent with our proposal. On the other hand, if the white dwarfs are real and in the halo, a young population is even more difficult

to explain than an old one. Given this, and the difficulty of the age calculations (which also couple to the difficulty in determining yields accurately), we require that the Pop III white dwarf relics are old. However, further study of this issue is clearly warranted.

The recently reported observations of an intrinsic dispersion in high-redshift D, taken together with the observation of a new Galactic white dwarf population, demand a non-standard history of star formation and chemical evolution. If these observations persist, then an unconventional scenario of the kind we propose will be *required*. We note that standard scenarios of chemical evolution are unable to lead to significant dispersion at low metallicity. If these standard models prevail, then one must concur with Pettini & Bowen (2001) that the D dispersion at high redshift is an artifact, and that significant systematic errors are present. At present, the use of D/H as a precision baryometer must be called into question until these systematic errors are better understood. We expect that future analyses and observations will either show a convergence towards a unique observed D/H value, representative of the primordial one, or reveal the dependence of D/H on metallicity in high z absorbing clouds.

It is a pleasure to thank Robert Mochkovitch, Patrick Petitjean and Jim Truran for stimulating discussions. BDF acknowledges enlightening conversations with the participants of the “Deuterium in the Universe” workshop. The work of KAO was supported in part by DOE grant DE-FG02-94ER-40823 at the University of Minnesota. KAO also acknowledges support during a sabbatical at CERN where much of this work was done. This work has been supported in part by PICS 1076 from the CNRS.

REFERENCES

- Adams, F. C. & Laughlin, G. 1996, ApJ, 468, 586
- Alcock, C. *et al.* 2000, ApJ, 542, 281
- Bonifacio, P. & Molaro, P. 1997, MNRAS, 285, 847
- Burles, S., & Tytler, D. 1998a, ApJ 499,699
- Burles, S., & Tytler, D. 1998b, ApJ 507, 732
- Chabrier, G. 1999, ApJ, 513, L103
- Chieffi, A., Domínguez, I., Limongi, M., & Straniero, O. 2001, ApJ, in press (astro-ph/0103104)

- Cole, S. *et al.* 2001, astro-ph/0012429, MNRAS in press
- Cyburt, R.H., Fields, B.D., & Olive, K.A. 2001, New Astronomy, 6, 215
- D’Odorico, S., Dessauges-Zavadsky, M., & Molaro, P. 2001, A&A, 368, L21
- Fields, B. D., Freese, K., & Graff, D. S. 2000, ApJ, 534, 265
- Fields, B. D. & Olive, K. A. 1999, New Ast, 4, 255
- Fujimoto, M. Y., Ikeda, Y., & Iben, I. 2000, ApJ, 529, 25
- Fukugita, M., Hogan, C., & Peebles, P.E.J. 1998, ApJ, 503, 518
- Gates, E. I. & Gyuk, G. 2001, ApJ, 547, 786
- Gautier, D., & Morel, P. 1997, A&A, 323, L9
- Geiss, J., & Gloeckler, G. 1998, Space Sci. Rev., 84, 239
- Hansen, B. 2001, astro-ph/0105018, ApJL, submitted.
- Hauser, M. and Dwek, E. 2001, astro-ph/0105363, ARAA, in press.
- Ibata, R., Irwin, M., Bienaymé, O., Scholz, R., & Guibert, J. 2000, ApJ, 532, L41
- Kirkman, D., Tytler, D., Burles, S., Lubin, D., O’Meara, J. M. 1999, ApJ, 529, 655
- Kirkman, D. *et al.* , 2001, astro-ph/0103305
- Kobayashi, C. *et al.* 1998, ApJ, 503, L155
- Koopmans, L.V.E., & Blandford, R.D. 2001, The Shapes of Galaxies and their Halos, ed. P. Natarajan, in press (astro-ph/0106392)
- Lauroesch, J.T., Truran, J.W., Welty, D.E., & York, D.G. 1996, PASP, 108, 641
- Linsky, J., 1998, Space Sci. Rev., 84, 285
- Marigo, P., Girardi, L., Chiosi, C., & Wood, P. R. 2001, A&A, 371, 152
- Matteucci, F. & Greggio, L. 1986, AA, 254, 279
- Molaro, P., *et al.* 2000, ApJ, 541, 54
- Netterfield, C.B. *et al.* 2001, astro-ph/0104460

- Nollett, K.M. & Burles, S. 2000, Phys. Rev. D, 61, 123505
- Olive, K.A. & Skillman, E. 2001, New Astronomy, 6, 119
- O’Meara, J.M., *et al.* 2001, ApJ, 522, 718
- Oppenheimer, B.R., Hambly, N.C., Digby, A.P., Hodgkin, S.T., & Saumon, D. 2001, Science, 292, 698
- Péroux, C., McMahon, M. and Storrie-Lombardi, L. and Irwin, M. 2001, astro-ph/0107045, MNRAS, submitted.
- Pettini, M., & Bowen, D.V. 2001, (astro-ph/0104474)
- Pettini, M., Ellison, S.L., Stidel, C.C., Shapley, A.E., & Bowen, D.V. 2000, ApJ, 532, 65
- Prantzos, N., & Ishimaru, Y. 2001, A&A, in press (astro-ph/0106497)
- Pinsonneault, M.H., Walker, T.P., Steigman, G., & Narayanan, V.K. 1998, ApJ, 527, 180
- Pinsonneault, M.H., Steigman, G., Walker, T.P., & Narayanan, V.K. 2001, astro-ph/0105439
- Prochaska, J.X., & Wolfe, A.M. 1999, ApJS, 121, 369
- Pryke, C., Halverson, N., Leitch, E., Kovac, J., Carlstrom, J., Holzappel, W. and Dragovan, M. 2001, astro-ph/0104490, ApJ in press.
- Riedeger, R., Petitjean, P. and Mucket, J.P., 1998, AA 360, 15
- Reid, N., Sahu, K. and Hawley 2001, astro-ph/0101110, ApJ in press.
- Ruiz-Lapuente, P., Cassé, M., & Vangioni-Flam, E. 2001, ApJ, 549, 483
- Ryan, S.G., Beers, T.C., Olive, K.A., Fields, B.D., & Norris, J. 2000, ApJ, 530, L57
- Ryan, S.G., Norris, J., & Beers, T.C. 1999, ApJ, 523, 654
- Ryu, D., Olive, K. A., & Silk, J. 1990, ApJ, 353, 81
- Smecker, T. and Wyse, R. 1991, ApJ, 372, 448
- Theado, S. & Vauclair, S. 2001, astro-ph/0106080
- Tytler, D., et al. 1999, AJ, 117, 63

- van den Hoek, L. B., & Groenewegen, M. A. T. 1997, A&AS, 123, 305
- Vangioni-Flam, E., Cassé, M., Cayrel, R., Audouze, J., Spite, M., & Spite, F. 1999, New Ast, 4, 245
- Vangioni-Flam, E., Coc, A., & Cassé, M. 2000, A&A, 360, 15
- Vauclair, S., & Charbonnel, C. 1998, ApJ, 502, 372
- Vidal-Madjar, A. 2001, in "Cosmic evolution", proceedings in the 60th birthdays of Jean Audouze and Jim Truran, World Scientific, Edts, E. Vangioni-Flam, R. Ferlet and M. Lemoine, in press (astro-ph/0103170)
- Wadsley, J. Hogan, C. and Anderson, S. 2000, in "Clustering at high redshift" in ASP conference series, Edts, A. Mazure, O Le Fevre and V. Le Brun, vol. 200, p. 290
- Webb, J.K., et al. 1997, Nature, 388, 250